



- HOME
- ABOUT
- RISK FACTORS
 - High Blood Pressure
 - Diabetes
 - Atrial Fibrillation
 - High Cholesterol
 - Physical Inactivity
 - Poor Diet
 - Alcohol
 - Obesity
 - Smoking
 - Stress and Depression
- RESOURCES
 - Videos
 - Blog
 - Podcast
 - Useful links
 - Additional Reading
 - Media Releases

- News
-  English
 -  English
 -  Português
 -  Русский
 -  Español
 -  Français
 -  Polski

- CONTACT

DONATE

10 Fascinating Facts About Stroke and High Blood Pressure



Ten fascinating facts about high blood pressure and stroke

Did you know that weakened arteries in the brain, resulting from high blood pressure, increase the risk for stroke? Here are ten more fascinating facts that

link high blood pressure and stroke and outline the importance of keeping blood pressure under control.

1. More than one billion people have high blood pressure worldwide

High blood pressure is more common than you might think. Over one billion people are estimated to suffer from the condition worldwide and, with often little or no symptoms, many more are undiagnosed (1).

2. High blood pressure figures have doubled since 1990

Lifestyle factors, such as physical inactivity, a salt-rich diet with highly processed and fatty foods, and alcohol and tobacco use, are at the centre of an increase in high blood pressure worldwide. This is spreading at an alarming rate from developed countries to low- or middle-income countries (2).

3. Heart disease and stroke are leading causes of death worldwide

The rise in the number of adults with high blood pressure is directly linked to the incidence of deaths from heart disease and stroke, two of the biggest killers in the world (3). Despite several initiatives, the prevalence of high blood pressure and cardiovascular deaths are increasing globally (4).

4. 90-95% of adult cases are classed as primary high blood pressure

There are two types of high blood pressure: primary and secondary. Primary, which may develop as a result of environmental or genetic causes, accounts for 90-95% of adult cases. While secondary, caused by other underlying conditions or medications, only accounts for up to 2-10% of cases (5).

5. People over 40 should have blood pressure checked regularly

Regular blood pressure monitoring is the key to preventing and managing high blood pressure. It is generally recommended that all adults over 40 years of age, without any other risk factors, have their blood pressure checked at least every five years so that any potential problems can be detected early (6).

6. Normal blood pressure should be lower than 120/80

Your blood pressure reading consists of two figures: systolic and diastolic. A normal reading should be lower than 120/80. High blood pressure is diagnosed if the first number (systolic pressure) is above 140 and the second number (diastolic pressure) is above 90 (4).

7. High blood pressure causes millions of deaths each year

High blood pressure affects around one billion people and causes an estimated 10.7 million deaths every year (2). But, known as the 'silent killer', around one in three adults aren't even aware they have it. That's why it's so important to get your blood pressure checked every year (3).

8. High blood pressure is more common in the over-65s

Many factors influence your likelihood of developing high blood pressure. Some are modifiable - for instance, you could stop drinking alcohol or smoking. Others are non-modifiable, such as old age: people over the age of 65 are more likely to suffer from it (4).

9. There are three main groups of blood pressure medications

The main types of blood pressure medication are thiazide and thiazide-like agents; angiotensin-converting enzyme (ACE) inhibitors/angiotensin-2 receptor blockers (ARBs); and long-acting calcium channel blockers. Your doctor may need to try different combinations to find out what works best for you (7).

10. It's easy to prevent high blood pressure and reduce stroke risk

There are several different things you can do to reduce your chance of stroke, caused by high blood pressure. These include losing weight; exercising more; limiting your tobacco, salt and alcohol intake; reducing stress; and getting a good night's sleep (4).

References

1. Tu N Nguyen, Clara K Chow. *Global and national high blood pressure burden and control. Lancet. 2021 August 24. Volume 398, issue 10304, P932-933. DOI: [https://doi.org/10.1016/S0140-6736\(21\)01688-3](https://doi.org/10.1016/S0140-6736(21)01688-3).*
2. *Worldwide trends in hypertension prevalence and progress in treatment and control from 1990 to 2019: a pooled analysis of 1, 201 population-representative studies with 104 million participants' by the NCD Risk Factor Collaboration (NCD-RisC). The Lancet.*
3. Virani SS, Alonso A, Benjamin EJ, Bittencourt MS, Callaway CW, Carson AP, et al. *Heart disease and stroke statistics-2020 update: a report from the American Heart Association external icon. Circulation 2020 and 141:e139-596. .*
4. Thomas Unger et. al. 2020 International Society of Hypertension. *Global Hypertension Practice Guidelines. 6 May 2020 <https://doi.org/10.1161/HYPERTENSIONAHA.120.15026>Hypertension.; 75:1334-1357 .*
5. Matthew R Alexander, MD, PhD et al. *Hypertension. Updated 2019 Feb 22. <https://emedicine.medscape.com/article/241381-overview#a4>.*
6. Walter N. Kernan, Silvio E. Inzucchi. *Treating Diabetes to prevent stroke. Originally published 20 Apr 2021. <https://www.ahajournals.org/doi/full/10.1161/STROKEAHA.120.032725>.*
7. *Organisation, Guideline for the pharmacological treatment of hypertension in adults. Geneva: World Health and 3.0.IGO, 2021. Licence CC By-NC-SA.*

Updated February 2022

Next review 2024



Developed by



SAFE retains full editorial control over the content of this website.

Supported by an educational grant from



Links

- [Terms Of Use](#)
- [Privacy Policy](#)
- [Cookie Policy](#)
- [Contact](#)

- [Facebook](#)
- [X](#)
- [RSS](#)